

PCT/EP 10/580777
Rec'd PCT/PTO 19 JUL 2004
INVESTOR IN PEOPLE

PRIORITY DOCUMENT

ITTED OR TRANSMITTED IN
ANCE WITH RULE 17.1(a) OR (b)

REC'D 25 APR 2003

WIPO

PCT

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.


In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed *Andrew Gensey*

Dated 13 December 2002

BEST AVAILABLE COPY



1/77

Request for grant of a patent

See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

17JAN02 E688535-1 CT4268
P01/7700 0.00-0200977.7

Cardiff Road
Newport
South Wales
NP10 8QQ

Your reference

L.B.P. 27 GB

1. Patent application number
(The Patent Office will fill in this part)

0200977.7

17 JAN 2002

2. Full name, address and postcode of the or of each applicant (underline all surnames)

Lonza Biologics plc.

228 Bath Road, Slough SL1 4DY

Berkshire/Great Britain

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

United Kingdom

3. Title of the invention

Glutamine-auxotrophic human cells capable of producing proteins and capable of growing in a glutamine-free medium

4. Name of your agent (if you have one)

Deana O Reilly

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Lonza Biologics plc.

Legal Office

228 Bath Road

Slough, Berkshire SL1 4DY / Great Britain

Patents ADP number (if you know it)

5. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

YES

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	1
Description	24
Claim(s)	9/2
Abstract	1
Drawing(s)	4 + 4 + 4

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11. I/We request the grant of a patent on the basis of this application

Signature Dr. John Birch

Date Nov. 23,

12. Name and daytime telephone number of person to contact in the United Kingdom

Deana O Reilly
+44 1753 716590

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 50051
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

The present invention relates to a novel glutamine-auxotrophic human cell capable of producing a protein and capable of growing in a glutamine-free medium. Furthermore, it relates to a novel process of producing a protein and to the use of a glutamine synthetase (GS) as a selectable marker in glutamine-auxotrophic human cells.

Protein production by mammalian cell culture is used to provide proteins for therapeutic and diagnostic applications. Today mammalian cell cultures are the preferred source of a number of important proteins for use in human and animal medicine, especially those which are relatively large, complex and glycosylated (N. B. Finter et al., in Large-Scale Mammalian Cell Culture Technology, 1990, ed. A. S. Lubiniecki, Marcel Dekker, Inc., New York).

For example, production of the human protein erythropoietin (EPO) by cell culture has been described in WO 93/09222. Human EPO has been obtained at appreciable specific production rates employing human fibroblast cells transfected with the exogenous gene coding for human EPO. In a further process for the production of human EPO (WO 94/12650), the human fibrosarcoma cell line HT1080 transfected with DNA sequences which are capable to activate the endogenous encoded EPO has been applied. A similar approach has been described in WO 99/09268. Immortalised human cells like Namalwa, Hela S3 and HT 1080 cells have been transfected with DNA sequences which are capable to activate the endogenous encoded EPO. The cells described in WO 93/09222, WO 94/12650 and WO 99/09268 used for the production of human EPO have all been cultured in a medium containing glutamine. This is disadvantageous because when glutamine is used as an energy substrate by cultured cells, ammonia is a catabolite produced, which is cytotoxic and can inhibit cell growth. Furthermore, it can inhibit glycosylation of proteins by its effect on pH within the Golgi of the cell.

The production of high levels of tissue plasminogen activator, a glycosylated protein, has been described in WO 87/04462. GS has been used herein as amplification system for co-amplifying the gene encoding tissue plasminogen activator (tPA) in glutamine-prototrophic Chinese hamster ovary (CHO) cells by transfecting the cells with a gene encoding GS. As found in EP-A 148 605, it is disadvantageous, however, to use CHO cells for the production of a glycosylated human protein. Proteins synthesized by CHO cells may differ in their average carbohydrate composition from natural occurring glycosylated human proteins. This is due to the fact that human cells possess $\alpha 2.3$ sialyltransferase and $\alpha 2.6$ sialyltransferase enzymes. CHO cells possess only the $\alpha 2.3$ sialyltransferase and so cannot perform the $\alpha 2.6$

linkage of terminal sialic acid to the oligosaccharide moieties. CHO cells lack the enzymes for sulphation of the carbohydrate structures. CHO cells lack also an α 1-3 fucosyltransferase (attaches terminal fucose residues) though do have α 1-6 fucosyltransferase (attaches core fucose residues). Human cells have both fucosyltransferases. (Cumming D.A., 1991, Glycobiology Vol. 1, No. 2, 115 - 130, Jenkins N. and Curling E.M.A., 1994, Enzyme and Microbial Technology Vol. 16, 354-364. Lee et al., 1989, Journal of Biological Chemistry, Vol. 264, 13848-13855). Therefore, glycosylated proteins synthesized by CHO cells may not have the desired characteristics e. g. in-vivo biological activity as the one produced in human cells.

In WO 89/10404, a method of making myeloma cells such as mouse hybridoma, mouse plasmacytoma cells and rat hybridoma cells glutamine-independent by transforming them with GS has been reported. It has been further demonstrated herein that GS can be used for co-amplification of genes encoding light and heavy chains of immunoglobulin molecules and for co-amplification of a gene encoding a fibrinolytic enzyme in a myeloma cell line. However, rodent cell lines present disadvantages, namely the attachment of N-glycolylneuraminic acid residues in place of the N-acetylneuraminic acid, the inability to carry out sulphation and the presence of the α 1.3 galactosyltransferase enzyme. Oligosaccharide structures of glycoproteins synthesized in rodent cells might therefore be expected to be immunogenic in humans.

The object of the present invention is to provide an improved process which does not have the above-mentioned disadvantages for the production of a protein, especially for the production of a glycoprotein, and which yields high protein titres.

This object has been achieved with a novel glutamine-auxotrophic human cell according to claim 1 and with a novel process according to claim 7.

According to the invention a glutamine-auxotrophic human cell is provided which has been transfected with an exogenous DNA sequence encoding a protein or an exogenous DNA sequence capable of altering the expression of an endogenous gene encoding a protein and an exogenous DNA sequence encoding a glutamine synthetase, wherein these exogenous DNA sequences are located on one or more than one DNA construct, said transfected cell capable of producing said protein and capable of growing in a glutamine-free medium.

Figure 1 shows adaptation of cell line R223 to suspension culture in serum-free medium - profiles of cell concentration during repeated serial subculture.

Figure 2 shows schematic for adaptation of HT1080 cells to suspension culture in serum-free medium.

Figure 3 shows adaptation of cell line HT1080 to suspension culture in serum-free medium - profiles of cell concentration during repeated serial subculture.

Figure 4 shows IEF analysis of immunopurified EPO from GS transfectant 3E10 and from the non-transfected R223 cell line. Lane 2: 3E10 harvest. Lane 3: 3E10 peak. Lane 4: non-transfected R223 cell line peak. Lane 5: non-transfected R223 cell line harvest.

DNA sequences

An "exogenous DNA sequence encoding a protein" or an "exogenous DNA sequence capable of altering the expression of an endogenous gene encoding a protein" and an "exogenous DNA sequence encoding a GS" are usually located on a DNA construct like an expression vector or an infectious vector. As expression vector a plasmid can be employed. As infectious vectors, can be employed e.g. retroviral, herpes, adenovirus, adenovirus-associated, mumps and poliovirus vector. Preferably an expression vector, in particular a plasmid is used.

An "exogenous DNA sequence encoding a protein" may include additional sequences such as a regulatory sequence e.g. a promoter and/or an enhancer, polyadenylation sites and splice junctions usually employed for the expression of the exogenous gene or may include additionally one or more separate targeting sequences and optionally DNA encoding a selectable marker as described in WO 93/09222.

An "exogenous DNA sequence capable of altering the expression of the endogenous gene encoding a protein" may include exogenous DNA sequences which do not encode a gene product of the protein but encode part of that gene product e. g. an exon, and may include additional sequences such as regulatory sequences and splice junctions usually employed for

the expression of the exogenous DNA sequence. They may further include targeting sequences and optionally DNA encoding a selectable marker as described in WO 93/09222. Usually, an "exogenous DNA sequence capable of altering the expression of an endogenous gene encoding a protein" is inserted into chromosomal DNA of the cell after transfection into the cell. Homologous recombination or targeting is hereby used to replace or disable the regulatory region normally associated with the endogenous gene with a regulatory sequence. As regulatory sequence serve e. g. a promoter and / or an enhancer which causes the gene to be expressed at levels higher than evident in the corresponding nontransfected cell as described e. g. in WO 93/09222. Appropriate promoters can be regulatable or constitutively expressed promoters. Appropriate promoters may be strong promoters which are depending on the cell line used e.g. the human cytomegalovirus major immediate early promoter (hCMV-MIE), SV 40 early and late promoters, other promoters of the adenoviruses, early and late promoters of any of the polyoma viruses or papova-viruses, interferon α 1 promoter, mouse metallothionein promoter, the rous sarcoma virus long terminal repeat promoter, β -globin promoter, conalbumin promoter, ovalbumin promoter, mouse β -globin promoter and human β -globin promoter.

The "exogenous DNA sequence encoding a GS" might be under the control of a strong promoter as well as under the control of a weak promoter. A strong promoter is used if the exogenous DNA sequence is required simply to express the gene encoding GS. A weak promoter is used if the exogenous DNA sequence is being used as a selectable marker and if GS is used for amplification. An appropriate promoter can be a regulatable or a constitutively expressed promoter. The promoter might be selected such as, that the GS is expressed at a concentration sufficient for growth of the transfected cell but which does not produce a high level of the glutamine catabolite product ammonia in cell culture, usually not more than 4 mM, preferably not more than 2 mM, more preferably less than 2 mM ammonia.

A "selectable marker" confers a selectable phenotype which makes it possible to identify and isolate recipient cells. GS can be used as the selectable marker in the present invention in order to select successfully transfected glutamine-auxotrophic human cells which have incorporated and express the exogenous DNA sequence encoding GS.

A strong promoter may, depending on the cell line used, be e. g., hCMV-MIE, SV 40 early and late promoters, other promoters of the adenoviruses, early and late promoters of any of the polyoma viruses or papova-viruses, interferon α 1 promoter, mouse metallothionein promoter,

the rous sarcoma virus long terminal repeat promoter, β -globin promoter, conalbumin promoter, ovalbumin promoter mouse β -globin promoter and human β -globin promoter. A weak promoter may, depending on the cell line used, be e.g. murine leukaemia virus long terminal repeat, herpes simplex virus thymidine kinase and Mouse Mammary Tumor Virus-Long Terminal Repeat. Preferably the gene encoding a GS is under the control of a strong promoter, more preferably under the control of the hCMV-MIE promoter.

The "exogenous DNA sequence encoding a protein" or the "exogenous DNA sequence capable of altering the expression of an endogenous gene encoding a protein" and the "exogenous DNA sequence encoding a GS" are located on one or more than one DNA construct. Preferably, these exogenous DNA sequences are located on more than one, more preferably on two DNA constructs. If these exogenous DNA sequences are located on one DNA construct they might be functionally combined, e.g. in that their expression is driven by the same regulatory sequence e.g. promoter and/or enhancer as described e.g. in WO 89/10404.

Human cells

Glutamine-auxotrophic human cells means all human cells which do not express GS or express GS poorly, thus being capable of growth in a culture medium containing glutamine but failing to grow or growing only poorly in glutamine-free medium. Glutamine-auxotrophic human cells which are used in the present invention are mortal glutamine-auxotrophic human cells or immortalized glutamine-auxotrophic human cells. Mortal glutamine-auxotrophic human cells are glutamine-auxotrophic human cells which exhibit a limited lifespan in culture. Immortalized also called "permanent" or "established" glutamine-auxotrophic human cells are glutamine-auxotrophic cells which exhibit an apparently unlimited lifespan in culture.

Examples of mortal glutamine-auxotrophic human cells may be human fibroblasts and human foetal lung tissue cells. Examples of immortalized glutamine-auxotrophic human cells may be human fibrosarcoma cells, like a HT1080 cell line and B lymphoblastoid human cells like a HL60 cell line. Preferably used in the present invention are immortalized glutamine-auxotrophic human cells. More preferably human fibrosarcoma cells, in particular a HT1080 cell line (ATCC No. CCL 121) is used.

Transfection

The glutamine-auxotrophic human cell can be transfected with the exogenous DNA sequences by known genetic engineering techniques.

Transfection with the exogenous DNA sequences depends on whether the sequences are

- 5 located on one or more than one DNA construct. If the sequences are located on more than one DNA construct, transfection can occur with each sequence separately or by co-transfection. In case transfection occurs with each sequence separately the order of transfection of the sequences is usually optional. Transfection with each sequence separately occurs preferably firstly with the "exogenous DNA sequence encoding said protein" or the
- 10 "exogenous DNA sequence capable of altering the expression of an endogenous gene encoding said protein" and secondly with the "exogenous DNA sequence encoding a GS". The transfected glutamine-auxotrophic cell might be cultured after each separate transfection and assessed for protein production.

- 15 In order to select for successfully transfected cells, these are grown in a glutamine-free medium. Cells might be grown directly in a glutamine-free medium or at first in a medium containing glutamine which will be diluted stepwise to a glutamine-free medium, e.g. one may start with a glutamine concentration of 10 mM which may be diluted by steps of 2 mM to 0 mM. The appropriate selection procedure might be chosen depending on the cell lines used.
- 20 As evident from the aforesaid, the glutamine-auxotrophic human cell of the present invention capable of producing a protein and capable of growing in a glutamine-free medium is obtainable by transfecting said cell with an exogenous DNA sequence encoding said protein or an exogenous DNA sequence capable of altering the expression of an endogenous gene encoding said protein and an exogenous DNA sequence encoding a glutamine synthetase,
- 25 wherein these exogenous DNA sequences are located on one or more than one DNA construct.

Amplification

- The exogenous DNA sequence encoding a protein or the exogenous sequence capable of
- 30 altering the expression of an endogenous gene encoding a protein may be amplified after transfection according to known methods in gene amplification as described in e.g. WO 94/12650. Amplifiable genes encoding enzymes like e.g. DHFR (dihydrofolate reductase), GS, adenosine deaminase, asparagine synthetase, aspartate transcarbamylase, metallothionein-

l, ornithine decarboxylase, P-glycoprotein, ribonucleotide reductase, thymidine kinase or xanthine-guanine phosphoribosyl transferase can be used for this purpose. Cells containing amplified copies of these genes are e.g. capable of surviving treatment in media lacking the metabolic product of the enzymes or in media containing a corresponding selective agent. Corresponding selective agents are e.g. methotrexate (MTX) in case of DHFR and methionine sulphoximine (MSX) in the case of GS.

Proteins

Proteins which are produced by the transfected glutamine-auxotrophic human cell of the present invention are non-glycosylated and glycosylated proteins. Glycosylated proteins refer to proteins having at least one oligosaccharide chain.

Examples for non-glycosylated proteins are e. g. non-glycosylated hormones like luteinizing hormone-releasing hormone, thyroid hormone-releasing hormone, insulin, somatostatin, prolactin, adrenocorticotrophic hormone, melanocyte-stimulating hormone, vasopressin, and derivatives thereof e. g., desmopressin, oxytocin, calcitonin, parathyroid hormone (PTH) or fragment thereof (e. g. PTH (1-43)), gastrin, secretin, pancreaticozym, cholecystokinin, angiotensin, human placental lactogen, human chorionic gonadotropin (HCG), caerulein and motilin; non-glycosylated analgesic substances like enkephalin and derivatives thereof (see US-A 4 277 394 and EP-A 031567), endorphin, dynorphin and kyotorphin; non-glycosylated enzymes like non-glycosylated nerve transmitters e. g. bombesin, neurotensin, bradykinin and substance P; non-glycosylated growth factors of the nerve growth factor (NGF) family, of the epithelial growth factor (EGF) and of the fibroblast growth factor (FGF) family and non-glycosylated receptors for hormones and growth factors.

Examples for glycosylated proteins are hormones and hormone releasing factors like growth hormones, including human growth hormone, bovine growth hormone, growth hormone releasing factor, parathyroid hormone, thyroid stimulating hormone, EPO, lipoproteins, alpha-1-antitrypsin, follicle stimulating hormone, calcitonin, luteinizing hormone, glucagon, clotting factors such as factor VIIIc, factor IX, tissue factor, and von Willebrands factor, anti-clotting factors such as Protein C, atrial natriuretic factor, lung surfactant, a plasminogen activator, such as urokinase or human urine or tissue-type plasminogen activator (t-PA), thrombin, hemopoietic growth factor, enkephalinase, RANTES (regulated on activation normally T-cell expressed and secreted), human macrophage inflammatory protein (MIP-1-alpha), a serum albumin such as human serum albumin, mullerian-inhibiting substance, relaxin A-chain,

relaxin B-chain, prorelaxin, mouse gonadotropin-associated peptide, a microbial protein, such as beta-lactanase, DNase, inhibin, activin, renin, vascular endothelial growth factor (VEGF), receptors for hormones or growth factors, integrin, protein A or D, rheumatoid factors, a neurotrophic factor such as bone-derived neurotrophic factor (BDNF), neurotrophin-3, -4, -5 or -6 (NT-3, NT-4, NT-5 or NT-6), or a nerve growth factor such as NGF- β , platelet-derived growth factor (PDGF), fibroblast growth factor such as a FGF and bFGF, epidermal growth factor (EGF), transforming growth factor (TGF) such as TGF- α and TGF- β , including TGF- β 1, TGF- β 2, TGF- β 3, TGF- β 4 or TGF- β 5, insulin-like growth factor-I and -II (IGF-I and IGF-II), des(1-3)-IGF-I (brain IGF-I), insulin-like growth factor binding proteins, CD proteins (cluster of differentiation proteins) such as CD-3, CD-4, CD-8 and CD-19, osteoinductive factors, immunotoxins, a bone morphogenetic protein (BMP), cytokines and their receptors, as well as chimeric proteins comprising cytokines of their receptors, including, for instance tumor necrosis factor α and β , their receptors (TNFR-1, EP 417 563, and TNFR-2, EP 417 014) and their derivatives, an interferon such as interferon- α , - β and - γ , colony stimulating factors (CSFs), e.g., M-CSF, GM-CSF and G-CSF, interleukins (ILs), e.g., IL-1 to IL-10, superoxide dismutase, T-cell receptors, surface membrane proteins, decay accelerating factor, viral antigen such as, for example, a portion of the AIDS envelope, transport proteins, homing receptors, addressins, regulatory proteins, antibodies, chimeric proteins, such as immunoadhesins, and fragments of any of the above listed glycosylated proteins. Preferably, glycosylated proteins are produced in the present invention. Most preferably glycosylated hormones like EPO, in particular EPO is produced.

Cell Culture

Any suitable culture procedure and culture apparatus known in the art may be used for growing the transfected human cell of the present invention. As culture medium common glutamine-free basal medium supplemented with about 0.1 to 20 %, preferably 0.5 to 15 % serum as well as serum-free, glutamine-free common basal medium can be used. Also, common glutamine-free basal medium free of protein of animal origin can be used. Preferably serum-free glutamine-free common basal medium is used.

Sera that may be used are e. g. foetal bovine serum or adult bovine serum. Preferably foetal bovine serum is used. Common glutamine-free basal medium that may be used are e. g. glutamine-free Eagle's minimal essential medium (MEM) medium, glutamine-free Dulbecco's Modification of Eagle's Medium (DMEM), glutamine-free Iscove's DMEM

medium (N. Iscove and F. Melchers, Journal of Experimental Methods, 1978, 147, 923), glutamine-free Ham's F12 medium (R.G. Ham, Proceedings of National Academy of Science, 1965, 53, 288), glutamine-free L-15 medium (A. Leibovitz, American Journal of Hygiene, 1963, 78, 173), glutamine-free RPMI 1640 medium (G.E. Morre et al., The Journal of the American Medical Association, 1967; 199, 519), a glutamine-free proprietary medium and suitable ratio mixtures thereof.

Common supplements might be added to the common glutamine-free basal medium. Supplements that might be usually added contain proteins usually present in serum and optionally further ingredients which have a positive effect on cell growth and/or cell viability. Proteins usually present in serum are e.g. bovine serum albumin (BSA), transferrin and/or insulin. Further ingredients which have a positive effect on cell growth and/or cell viability are e.g. soybean lipid, selenium and ethanolamine. Amino acids which replace glutamine and/or nucleosides might be added to the culture medium depending on the cell line. Examples of amino acids are isoleucine, leucine, valine, lysine, asparagine, aspartic acid, glutamic acid, serine, alanine. Optionally glutamine might be added to the common glutamine-free basal medium at low concentrations of usually less than 1 mg/l, preferably less than 0.5 mg/l to support its biosynthetic function (e.g. transamination reactions).

If the exogenous DNA sequence encoding a protein or the exogenous sequence capable of altering the expression of an endogenous gene encoding a protein has been amplified after transfection using amplifiable genes, the corresponding selective agent may be added to the common glutamine-free basal medium. The applied concentration range of the selective agent does depend on the cell line used. Usually, concentrations of 10 μ M and higher are used.

Adaptation for growing in suspension in serum-free medium

A glutamine-auxotrophic human cell which can be used as starting material for obtaining a transfected glutamine-auxotrophic human cell capable of producing a protein and capable of growing in a glutamine-free medium according to the invention might be anchorage-dependent or anchorage-independent. If an anchorage-dependent human cell, e. g. the HT1080 cell line (ATCC No. CCL 121) is used, it can be adapted to be a anchorage-independent HT1080 cell line capable of growing in suspension in serum-free medium which has not been described in the literature yet.

Adaptation might occur before or after transfection with the exogenous DNA sequence encoding a protein, or the exogenous DNA sequence capable of altering the expression of an endogenous gene encoding a protein and the exogenous DNA sequence encoding a GS.

Preferably, the cell is firstly transfected with the exogenous DNA sequence encoding a protein or the exogenous DNA sequence capable of altering the expression of an endogenous gene encoding a protein, and secondly adapted for growing in suspension in serum-free medium and then further transfected with the exogenous DNA sequence encoding a GS. If necessary the transfected cell might be again adapted for growing in suspension in serum-free, glutamine-free medium.

10 The transfected glutamine-auxotrophic human cell of the present invention might be anchorage-dependent or anchorage-independent and might be capable of growing in suspension in serum-free, glutamine-free medium. The preferred transfected glutamine-auxotrophic human cell is anchorage-independent and is capable of growing in suspension in serum-free, glutamine-free medium.

15 The adaptation for being an anchorage-independent cell capable of growing in suspension in serum-free medium can be achieved by adapting the cell in a first step to be anchorage-independent using serum-containing medium. This can be done by e. g. trypsinisation of the cells and subsequent agitation or releasing the cells by agitation. The cells are then adapted in a second step to serum-free medium by subsequent reduction of serum content. During
20 adaptation the amount of selective agent if used can be reduced in case it is inhibitory to growth of the cells. However, the cells might be as well adapted in a first step to grow in serum-free medium by subsequent reduction of serum content and in a second step adapted to be anchorage-independent cells capable of growing in suspension by e. g. trypsinisation of the cells and subsequent agitation or releasing the cells by agitation. Both steps might be as well
25 applied simultaneously.

Preferably the cells are adapted in a first step to be anchorage-independent cells using serum-containing medium by releasing the cells by agitation and then adapting them in a second step to serum-free medium by subsequent reduction of serum content.

30 As a basis for serum-containing medium culture medium as described above can be used. Selective agents as defined herein may be added to the culture medium. The applied concentration range of the selective agent does depend on the cell line used. Usually concentration of 10 μ M and higher are used. Serum-containing medium is usually

supplemented with about 0.1 to 20 % preferably 0.2 to 10 % most preferably 0.5 to 5 % serum. Sera that may be used are as mentioned above. Subsequent reduction of serum content might be obtained by reducing the serum content stepwise e.g. from 10 % to 1 % to 0 %.

Process for protein production

The transfected glutamine-auxotrophic human cell of the present invention is used in a process for the production of a protein by culturing said cell in a culture medium under conditions suitable for expression of said protein and recovering said protein. Proteins produced are as described above. As culture medium common glutamine-free basal media and common supplements as described above can be used. Suitable culture conditions are those conventionally used for in vitro cultivation of mammalian cells as described e. g. in WO 96/39488.

Protein recovery

Protein can be isolated from the cell culture by conventional separation techniques such as e.g. fractionation on immunoaffinity or ion-exchange columns; precipitation; reverse phase HPLC; chromatography; chromatofocusing; SDS-PAGE; gel filtration. One skilled in the art will appreciate that purification methods suitable for the polypeptide of interest may require modification to account for changes in the character of the polypeptide upon expression in recombinant cell culture.

Example 1**Provision of the Human Fibrosarcoma Cell Line HT1080 R223**

The anchorage-dependent human fibrosarcoma HT 1080 cell line capable of producing human EPO containing multiple copies of the human EPO gene has been obtained from

- 5 Transkaryotic Therapies, Inc. Cambridge, MA 02139 (US). The HT1080 cell line (ATTC No CCL 121) has obtained the capability for producing EPO by transfection with the DNA construct pREPO22 which is similar to the DNA construct pREPO18 described in WO 95/31 560 except that the DHFR gene is in the opposite orientation and that pREPO22 does contain approximately 600 base pairs less homologous sequence than pREPO18. This cell line is
- 10 further referred to as the R223 cell line.

Example 2**Adaptation of the R223 Cell Line to Growth in Suspension in Serum-Free Medium**

- Cells grown as attached cultures in static flasks were released by trypsinization, resuspended
- 15 into a proprietary glutamine-containing serum-free medium further referred to as "HM9" supplemented with 10 % dialysed foetal bovine serum (dFBS) and 500 nM MTX and incubated as shake flask cultures. Growth commenced after 6 days and the cells were subcultured into the same medium (Figure 1). Once a reliable pattern of growth had become established the serum content of the medium was reduced to 1 %. Again reliable growth was
- 20 allowed to become re-established before complete elimination of the serum supplement. Serum-supplemented and serum-free cultures were overgrown to assess productivity. Results are shown in Table 1 together with data from attached cultures. After adequate adaptation specific growth rate in suspension culture, even in the absence of serum, was equivalent to that in serum-supplemented attached culture.
- 25 Upon adaptation from attached culture to suspension culture in serum supplemented medium the specific rate of EPO synthesis decreased by 50 %, from 24 to 12 EU/10⁶ cells/h. However, this rate was substantially restored, to 18 EU/10⁶ cells/h, following adaptation to serum-free growth.

Table 1

Growth and productivity of the R223 cell line in attached culture and in suspension culture before and after adaptation to serum-free medium.

	maximum cells/mL $\times 10^{-6}$	specific growth rate h^{-1}	cumulative cell h/mL $\times 10^{-6}$	EPO EU/mL	q_{EPO} EU/ 10^6 cells/h
attached culture (DMEM + 10 % dFBS 500 nM MTX)	not applicable	0.0157	not applicable	not applicable	24
suspension culture (HM9 + 10 % dFBS 500 nM MTX)	1.4	0.0185	334	4199	12
suspension culture (HM9 (serum-free) 500 nM MTX)	1.1	0.0166	210	3600	18

Example 3**Adaptation of the HT1080 ATCC CCL121 to Growth in Suspension in Serum-Free Medium**

The HT1080 cell line ATCC CCL121 was obtained from the American Type Culture
 5 Collection (Rockville, Maryland, USA). Initially cells were grown in attached cultures in
 DMEM containing 10 % foetal bovine serum (FBS).

The steps to suspension and serum-free adaptation can be performed according to the
 schematic represented in Figure 2. To initiate suspension cultures, cells were released from
 attached cultures by trypsinization and the released cells were resuspended into HM9 plus 2 %
 10 FBS. These were then incubated as shaken cultures. Once suspension growth of the cells in
 HM9 plus 2 % FBS had become established the cultures were diluted with the same medium
 to make daughter cultures. Reliable suspension growth of the HT1080 cell line was
 established after 30 days in culture (Figure 3). Thereafter the serum content of the medium
 was reduced, and finally eliminated completely. The cells continued to grow in serial
 15 subculture in the absence of serum.

Example 4**Effect of Ammonia on Growth and Productivity of the R223 Cell Line**

Ammonia is a catabolite produced by cultured cells when glutamine is used as an energy
 20 substrate. It is cytotoxic and can inhibit cell growth. Furthermore, it can inhibit glycosylation
 of proteins by its effect on pH within the Golgi of the cell. R223 cells in flask cultures
 typically produce 5 mM ammonia in unfed cultures and 10 mM in fermenter cultures which
 receive a nutrient feed.

In an initial examination of the effects of ammonia, R223 cells were grown in shake flasks,
 25 either without added ammonia, or with ammonia added at 2, 5 or 10 mM. For each
 concentration of ammonia replicate cultures were set up at three different pH values, by
 varying the CO₂ content of the overlay gas. The primary aim here was to determine the extent
 of growth inhibition exerted by ammonia and to test whether reduced pH would overcome this
 growth inhibition.

30 While ammonia was found to inhibit cell growth (Table 2), reduced pH failed to alleviate the
 inhibitory effects of ammonia, though the elevated pCO₂ required to reduce the pH may itself
 have caused some growth inhibition.

Cultures were terminated after only 3 days. To continue any longer was not valid because accumulation of non-volatile acid catabolites caused the pH of all cultures to fall by several points of a pH unit.

Table 2

Effect of Ammonia on growth and productivity of R223 cells at different values of culture pH. The pH was adjusted by the content of CO₂ in the overlay gas.

pCO ₂ %	ammonia added mM	initial pH	maximum cells/ml x 10 ⁻⁶	specific growth rate h ⁻¹	EPO EU/ml	Q _{EPO} EU/10 ⁶ cells/h
5	0	7.26	0.9	0.020	1719	35
10	0	6.93	0.7	0.017	1580	40
15	0	6.75	0.4	0.012	1186	56
5	2	7.15	0.8	0.018	1796	38
10	2	6.88	0.7	0.017	1928	42
15	2	6.80	0.3	0.009	1404	78
5	5	7.18	0.6	0.018	2639	40
10	5	6.91	0.6	0.014	1663	51
15	5	6.75	0.3	0.016	1626	45
5	10	7.17	0.5	0.016	2222	43
10	10	6.90	0.6	0.014	1966	50
15	10	6.75	0.2	0.008	1891	86

In a subsequent experiment (Table 3) pH was varied by adjusting the NaHCO₃ content of the medium while maintaining the CO₂ content of the overlay gas at a non-inhibitory concentration. The range of pH tested was from pH 7.0 to 7.5. (It must be emphasized that in flask cultures medium pH cannot be controlled and falls substantially, even over the first two days of culture. The pH values stated are the initial pH of the each culture). Specific growth rate was maximal at the pH 7.5 (NaHCO₃ at 3 g/l) but at this pH both growth rate and maximum cell concentration were halved by the presence of 10 mM ammonia while the specific rate of EPO synthesis was reduced four fold (Table 3).

At pH 7.25 (NaHCO₃ at 1.5 g/l) specific growth rate was reduced, and the specific rate of EPO synthesis was increased, compared to the culture at pH 7.5. However growth rate was less

affected, and the specific rate of EPO synthesis was unaffected, by the presence of 10 mM ammonia.

At the lowest pH studied, pH 7.0 (NaHCO_3 at 0.75 g/l) specific growth rate was further reduced but again was less affected by ammonia than at pH 7.5. The specific rate of EPO

5 synthesis was increased in the presence of added ammonia (this may have been attributable to the reduction in growth rate).

Table 3

10 **Effect of ammonia on growth and productivity of R223 cells at different values of culture pH. The pH was adjusted by varying the content of NaHCO_3 in the medium.**

	initial pH	maximum cells/ml $\times 10^{-6}$	specific growth rate h^{-1}	EPO EU/ml	q_{EPO} EU/ 10^6 cells/h
NaHCO_3 3 g/l	7.49	1.3	0.030	951	16
NaHCO_3 3 g/l 10 mM ammonia	7.51	0.5	0.016	256	4
NaHCO_3 1.5 g/l	7.26	1.1	0.027	1151	24
NaHCO_3 1.5 g/l 10 mM ammonia	7.27	0.7	0.021	1191	26
NaHCO_3 0.75 g/l	7.02	0.9	0.020	995	26
NaHCO_3 0.75 g/l 10mM ammonia	7.01	0.4	0.016	1009	42

Example 5**Productivity of the R223 Cell Line in Fermenter Cultures at 5 Litre Scale**

For the R223 cell line three 5 litre fermentations were carried out, controlled at different values of pH between 6.95 and 7.15 (see Table 4). Each culture received a concentrated nutrient feed containing amino acids and glucose designed to maintain major consumed nutrients at sufficient concentration. Cultures grew well at pH 7.15 and at pH 7.05, but the culture at pH 6.95 failed to grow, probably due to the high concentration of CO₂ required to control at that pH.

Table 4

Productivity and metabolism of the R223 cell line in fermenter culture.

fermentation conditions	maximum cells/ml $\times 10^{-6}$	doubling time hours	cumulative cell hours/ml $\times 10^{-6}$	EPO EU/ml	q_{EPO} EU/ 10^6 cells/h	Final Ammonia mg/l
pH 7.15 NaHCO ₃ at 2 g/l	1.0	46	197	10795	58	163
pH 7.05 NaHCO ₃ at 2 g/l	1.0	51	195	11414	64	157
pH 6.95 NaHCO ₃ at 2 g/l	0.2	105	62	2995	70	not done

Example 6. Transfection with GS and productivity of GS transfectants in attached culture.

Cells used for the transfection were from a suspension-adapted serum-free stock of the R223 cell line of example 2. As these cells grow as large multicellular aggregates they were trypsinised to reduce them to substantially single-cell suspension.

For transfection an aliquot of approximately 10^7 of the single-suspension adapted R223 cell line (in phosphate buffered saline without calcium or magnesium) obtained in example 2 was mixed with 20 μ g of linearized DNA containing the GS gene (DNA sequence pCMGS Bam HI described in Bebbington et al., 1992, Biotechnology 10, 169-175) and subjected to

electroporation using a Biorad Gene Pulser (450 volts, 250 μ F). As a control an equivalent aliquot of cells was electroporated without addition of DNA.

Cells were diluted with HM9 medium without MTX containing 0 or 10 % dFBS and distributed into 96-well culture plates or into 25 cm² flasks and incubated at 35.5 to 37 °C.

- 5 HM9 medium initially contained 2mM glutamine but this was diluted to 0.5 mM after 1 day and then replaced after ten days with glutamine-free HM9 medium. Also at day 1 or day 10, MTX (500 nM) was re-introduced to the cultures.

For the cultures set up with the control cells, which were electroporated without added DNA, no growth was obtained.

- 10 For the cells which were electroporated with DNA, 15 GS transfectants were identified in six 96-well plates. GS transfectants were obtained both with, and without, MTX in the initial medium. Of the 15 GS transfectants seven were successfully expanded to flask cultures (Table 5). The remaining eight GS transfectants exhibited aberrant cell morphology, or grew poorly, and were abandoned.

- 15 For each of the initial seven GS transfectants isolated, a set of replicate static flask cultures was set up using 10 % serum-supplemented glutamine-free HM9 medium. At intervals of 1 to 4 days, cultures were sacrificed in order to count the cells and to measure the concentration of EPO by EPO ELISA. From a composite of these data, the specific rate of EPO synthesis could be estimated for each GS transfectants. Data are summarized in Table 5 and, for comparison,
20 data for the non-transfected R223 cell line are included. All GS transfectants exhibited elevated specific rates of EPO synthesis compared to the non-transfected R223 cell line. The best GS transfectant, 3E10, had a synthesis rate five to six fold higher than the non-transfected R223 cell line.

Table 5

Specific rates of EPO synthesis for GS transfectants of the R223 cell line and for the non-transfected R223 cell line. Data are derived from attached cultures grown in the presence of 10% dFBS. Methotrexate was omitted from the medium used for making the transfectants, but was reintroduced to the cultures at day 1 or 10 after transfection.

GS transfectant	day of addition of MTX (500 nM) to transfection plate	q _{EPO} (attached) EU/10 ⁶ cells / h
3B3	10	59
3E10	10	139
3E11	10	45
4D9	10	35
8F11	1	47
8F12	1	79
8G3	1	52
non-transfected R223 cell line assessed in attached culture	not applicable	24

Example 7**Adaptation of GS transfectants to suspension culture and serum-free medium.**

Of the seven GS transfectants obtained in Example 6, GS transfectant 3E10 and GS transfectant 8G3 were progressed into suspension culture.

Cells were grown as attached cultures in static flasks in glutamine-free HM9 medium supplemented with 10 % dFBS and 500nM MTX at 35.5 to 37 °C. The cells were released after 5 or 6 days by agitation, resuspended into the same medium, and incubated as shake flask cultures at the same temperature. Growth commenced after only two or three days. Cells were subcultured once in the same medium and then overgrown to assess productivity. Titres and product synthesis rate were at least two fold higher than for the non-transfected R223 cell line grown in 10 % dFBS and 500 nM MTX (Table 6).

Table 6

Growth and productivity of GS transfectants 3E10 and 8G3 in suspension culture in shake flasks. Glutamine-free HM9 medium contained 10 % dFBS and 500 nM MTX.

GS transfectant	Max. cells / ml $\times 10^6$	Max. EPO titre EU / ml	Q_{EPO} (suspension) EU / 10^6 cells / h
3E10	0.7	11113	57
8G3	1.2	10171	33
non-transfected R223 cell line assessed in suspension culture grown in 10% dFBS and 500 nM MTX	1.4	4199	12

For suspension cultures of both GS transfectants 3E10 and 8G3 the dFBS content of the glutamine-free HM9 medium was reduced stepwise from 10 % to 2 % to 1 % to 0.2 % to 0.1 % to 0 % for 3E10 and 10 % to 2 % to 1 % for 8G3, allowing reliable cell growth to become established at each dFBS concentration before further reduction. 8G3 was not further adapted than to 1 % dFBS.

Example 8

Productivity and metabolism of GS transfectant 3E10 in serum-free suspension culture.

The GS transfectant 3E10 adapted to serum-free growth in Example 7 was cultured in suspension culture in serum-free glutamine-free HM9 medium at 35.5 to 37 °C. For the GS transfectant 3E10 cell line 2 to 3 fold higher specific rates of EPO synthesis were obtained than for the non-transfected R223 cell line, which was cultured under the same conditions in HM9 medium containing glutamine (Table 7).

Table 7

Growth and productivity of GS transfectant 3E10 cells in serum-free suspension culture.

	maximum cells / ml $\times 10^{-6}$	EPO EU / ml	q_{EPO} EU / 10^6 / h
3E10 (glutamine-free HM9 medium)	1.1 – 1.2	9731 – 14234	40 – 67
R223 (HM9 medium with 6 mM glutamine)	1.0 – 1.6	3075 – 5818	13 – 24

The higher EPO-productivity of the GS transfected cell was accompanied by the reduction in the release of metabolic ammonia. Instead of the 5 mM ammonia typically produced in flask cultures of the non-transfected R223 cell line in HM9 medium containing 6 mM glutamine, the GS transfectant 3E10 produced only 1.8 mM ammonia in glutamine-free HM9 medium (Table 8).

Table 8

Reduction in ammonia synthesis in GS transfectant 3E10.

	Max. cells / ml $\times 10^{-6}$	Ammonia mM	q_{Ammonia} $\mu\text{Moles} / 10^6 \text{ cells} / \text{h}$
R223 (HM9 medium containing 6 mM glutamine)	0.9	4.8	30
	1.2	5.0	23
	1.0	5.5	27
3E10 (glutamine-free HM9 medium)	0.8	1.8	11

Example 9

Analysis of Product Quality.

EPO produced by GS transfectant 3E10 in flask cultures has been immunopurified and analysed for its distribution of glycoforms. Figure 4 shows isoelectric focusing (IEF) gel analysis of EPO obtained at peak cell concentration and harvest of a culture of 3E10 cells

grown in glutamine-free HM9 medium as described in Example 8. Comparable samples from the non-transfected R223 cell line grown in HM9 medium are included. EPO produced from GS transfectant 3E10 exhibited intensification of the more acidic isoforms compared to the non-transfected R223 cell line. Scanning of the IEF gels has allowed quantitation of the isoform distribution and calculation of a theoretical isoform relative activity (IRA).

Table 9

Analysis of isoform relative activity for EPO from non-transfected R223 cell line and GS transfectant 3E10.

	Band number	Specific Activity*	non-transfected R223		GS transfectant 3E10	
			Percentage of each isoform	Activity (percent x specific activity)	Percentage of each isoform	Activity (percent x specific activity)
more basic	1		5.2		2.5	
	2		5.2		3.2	
	3		6.3		3.6	
	4		11.4		6.0	
	5		10.5		6.2	
	6	0.071	9.2	0.7	7.5	0.5
	7	0.194	8.8	1.7	10.5	2.0
	8	0.273	8.7	2.4	11.6	3.2
	9	0.373	5.5	2.1	9.3	3.5
	10	0.658	4.4	2.9	9.5	6.3
	11	0.989	4.3	4.3	7.8	7.7
	12	0.999	4.5	4.5	6.1	6.1
	13	1.000	1.4	1.4	5.3	5.3
	14	0.796	0.2	0.2	1.7	1.4
	15				0.5	
	16					
more acidic	17					
	other minor bands		14.4		8.7	
			Sum = 100 %	Sum = IRA = 20.2	Sum = 100 %	Sum = IRA = 36.0

* Values for activity derived from EP-A 0 428 267

The data indicated considerable enhancement of the product quality in the GS transfectant 3E10 (Tables 9 and 10), with an IRA almost two fold higher than for the control culture of the non-transfected R223 cell line.

Also the hypothetical N-glycan charge "Z" was determined for the EPO from GS transfectant 3E10. "Z" was not determined for the non-transfected R223 cell line in Table 10. Nevertheless the "Z" values for GS transfectant 3E10 (273 at peak cell concentration and 265 at harvest) exceed values obtained in flask culture (183-228) for the non-transfected R223 cell line. "Z" was determined according to Hermentin et al., *Glycobiology*, 1996, 6, 217 - 230.

Table 10

Analysis of product quality for EPO from non-transfected R223 cell line and GS transfectant 3E10.

		titre EU / mL	Isoform relative activity	'Z'
R223 parent (HM9 medium with 6mM glutamine)	peak	3518	25.2	not done
	harvest	6150	20.2	not done
3E10 (glutamine-free HM9 medium)	peak	3921	43.6	273
	harvest	6862	36.0	265

Example 10

Productivity, Metabolism and Product Quality of GS Transfectant 3E10 in Serum-Free Suspension Culture in 6 Litre Fermenter

The GS-transfectant 3E10 was grown in batch culture in airlift fermenters. The culture medium was glutamine-free HM9, without serum. Each culture received a concentrated nutrient feed, containing amino acids and glucose designed to maintain major consumed nutrients at sufficient concentration. Results are shown in Table 11 and data for a fermentation of the non-transfected parent R223 are included for comparison.

The GS transfectant exhibited extended viability, hence an increase in the duration of culture. The specific rate of product synthesis was equivalent to that of the non-transfected parent cell line unchanged, but the greater culture longevity resulted in increased maximum product

concentration. Ammonia accumulation was at least four fold lower for the GS transfectant. Product quality, measured by IRA, was improved for the GS transfectant.

Table 11

- 5 **Growth and productivity of GS transfectant 3E10 cells in serum-free suspension culture. Product quality of EPO produced was quantified by IRA.**

Fermentation conditions	Max cells/ml $\times 10^6$	Max CCH*/ml $\times 10^6$	Harvest day	Ammonia mg/L	Harvest Titre EU/ml	q_{EPO} overall EU/ 10^6 cells/h	IRA at harvest
glutamine-free	1.03	306	21	41	20936	69	43.6
glutamine-free	1.13	360	24	20	20150	52	42.6
glutamine-free	0.94	319	24	15	18680	49	48.5
parent R223 in standard process with glutamine	1.11	216	13	165	12225	58	28

* Cumulative cells hours, the time integral of cell concentration

Claims:

A glutamine-auxotrophic human cell transfected with an exogenous DNA sequence encoding a protein or an exogenous DNA sequence capable of altering the expression of an endogenous gene encoding a protein and an exogenous DNA sequence encoding a glutamine synthetase, wherein these exogenous DNA sequences are located on one or more than one DNA construct, said transfected cell capable of producing said protein and capable of growing in a glutamine-free medium.

The glutamine-auxotrophic human cell of claim 1, wherein the exogenous DNA sequences are located on more than one DNA construct.

The glutamine-auxotrophic human cell of any of claim 1 or 2, wherein the glutamine-auxotrophic human cell is an immortalized glutamine-auxotrophic human cell.

The glutamine-auxotrophic human cell of claim 3, wherein the immortalized glutamine-auxotrophic human cell is a human fibrosarcoma cell.

The glutamine-auxotrophic human cell of claim 4, wherein the human fibrosarcoma cell is a HT1080 cell line.

The glutamine-auxotrophic human cell of claim 1 to 5, wherein the transfected cell is anchorage-independent and capable of growing in suspension in serum-free, glutamine-free medium.

A process for the production of a protein comprising

- a) a glutamine-auxotrophic human cell according to claim 1 is cultured in a culture medium under conditions suitable for expression of said protein and
- b) said protein is recovered.

The process of claim 7 wherein the protein is a glycosylated protein.

9. The use of a glutamine synthetase as a selectable marker in glutamine-auxotrophic human cells.

Abstract of the disclosure:

A glutamine-auxotrophic human cell transfected with an exogenous DNA sequence encoding a protein or an exogenous DNA sequence capable of altering the expression of an endogenous gene encoding a protein and an exogenous DNA sequence encoding a glutamine synthetase, wherein these exogenous DNA sequences are located on one or more than one DNA construct, said transfected cell capable of producing said protein and capable of growing in a glutamine-free medium.

Figure 1




Figure 2




Figure 3





Figure 4



BEST AVAILABLE COPY